

Neutrino flux in the rotating reference frame

D.L. Khokhlov

Sumy State University, R.-Korsakov St. 2,

Sumy 40007, Ukraine

E-mail: khokhlov@cafe.sumy.ua

Abstract

It is considered neutrino flux in the rotating reference frame. Due to the rotation of the frame, neutrino is observed as a superposition of two states P-transformed one from another. Since P-transformation is forbidden for neutrino, in the rotating reference frame one can detect a half of neutrino flux. Due to the rotation of the earth, the detector of neutrinos can measure a half of the solar neutrino flux predicted by the SSM that may provide a solution for the solar neutrino puzzle.

Weak interactions violate P-invariance and C-invariance but conserve CP-invariance [1]. The operations of C-transformation and of P-transformation are forbidden for neutrinos, but the operation of CP-transformation is allowed for neutrinos.

Take the rotating reference frame. Turnover of the frame at the angle π corresponds to P-transformation of the frame. Under rotation, the positions of the frame in the range from π to 2π are P-transformed from the positions of the frame in the range from 0 to π . Neutrino in the rotating frame is observed in the superpositional state

$$|\psi\rangle = \frac{1}{\sqrt{2}}(|\nu\rangle + P|\nu\rangle). \quad (1)$$

However P-transformation is forbidden for neutrino therefore it is forbidden to detect neutrino in the state $P|\nu\rangle$. Hence when measuring neutrino flux in the rotating frame it is forbidden to detect a half of neutrino flux.

As known [2], [3] experiments measured solar neutrino fluxes significantly smaller than those predicted by the standard solar model (SSM), e.g. [4], [5]. Discrepancy between the theoretical and experimental neutrino flux from the sun presents the solar neutrino puzzle.

The Homestake experiment measured [6] an average ^{37}Ar production rate by solar neutrinos in ^{37}Cl of

$$\text{Rate (Chlorine)} = 2.55 \pm 0.17(\text{stat}) \pm 0.18(\text{syst}) \text{ SNU} \quad (2)$$

which is $32 \pm 5\%$ of the $8.1_{-1.2}^{+1.0} \text{ SNU}$ predicted by the SSM [5].

The Kamiokande II and III experiment reported [7] the data consistent with ^8B solar neutrino flux of

$$\phi_{\nu_{\odot}} = [2.9 \pm 0.2(\text{stat}) \pm 0.3(\text{syst})] \times 10^6 \text{ cm}^{-2} \text{ s}^{-1} \quad (3)$$

which is $51\% \pm 9\%$ of that predicted by the SSM [5].

The GALLEX experiment measured [8] a ^{71}Ge production rate by solar neutrinos in ^{71}Ga of

$$\text{Rate (Gallium)} = 79 \pm 10(\text{stat}) \pm 6(\text{syst}) \text{ SNU} \quad (4)$$

which is $60\% \pm 10\%$ of the 131.5^{+7}_{-6} SNU predicted by the SSM [5].

The SAGE experiment measured [9] an average ^{71}Ge production rate by solar neutrinos in ^{71}Ga of

$$\text{Rate (Gallium)} = 74^{+13}_{-12}(\text{stat})^{+5}_{-7}(\text{sys}) \text{ } SNU \quad (5)$$

which is $56\% \pm 11\%$ of the 131.5^{+7}_{-6} SNU predicted by the SSM [5].

Due to the rotation of the earth, the detector of neutrinos can measure a half of the solar neutrino flux predicted by the SSM. This is consistent with the data of the Kamiokande, GALLEX, SAGE experiments and significantly reduces the discrepancy between the theoretical and experimental data for the Homestake experiment. So the effect under consideration may provide a solution for the solar neutrino puzzle.

References

- [1] E.D. Commins, P.H. Bucksbaum, *Weak interactions of leptons and quarks* (Cambridge University Press, 1983)
- [2] J.N. Bahcall, Nucl. Phys. B (Proc. Suppl.) **38** (1995) 98
- [3] N. Hata et al., Phys. Rev. D **52** (1995) 3622
- [4] S. Turck-Chieze et al., Apj. **335** (1988) 415
- [5] J.N. Bahcall, & M. Pinsonneault, Rev. Mod. Phys. **64** (1992) 885
- [6] B.T. Cleveland et al., Nucl. Phys. B (Proc. Suppl.) **38** (1995) 47
- [7] Y. Suzuki, Nucl. Phys. B (Proc. Suppl.) **38** (1995) 54
- [8] P. Anselmann et al., Nucl. Phys. B (Proc. Suppl.) **38** (1995) 68
- [9] J.N. Abdurashitov et al., Nucl. Phys. B (Proc. Suppl.) **38** (1995) 60